International Journal of Scientific Research & Engineering Trends
Volume 1, Issue 4, July-2015, ISSN (Online): 2395-566X

A Comprehensive Review of Schiff Base Transition Metal Complexes: Synthetic Strategies, Structural Insights, and Biological Applications

[1]Ranjan Kumar, and [2]Niranjan Kumar Mandal
[1]Department of Chemistry, Godda College, Godda, India
[2]Department of Chemistry, S. K. M. University, Dumka, India

Abstract - Schiff base transition metal complexes have emerged as a pivotal class of compounds in coordination chemistry due to their structural diversity, facile synthesis, and broad-spectrum biological activities. This review critically compiles and analyzes literature from 2000 to 2013, emphasizing synthetic methodologies, coordination behavior, and characterization techniques of Schiff base ligands and their complexes with transition metals. Particular attention is given to their promising applications in antimicrobial, antioxidant, anticancer, and catalytic domains. The role of the azomethine (-C=N-) functional group in ligand-metal binding and its impact on physicochemical properties is discussed in detail. Advances in understanding structure-activity relationships highlight how metal ion identity and ligand substituents influence bioactivity and selectivity. Challenges such as limited substrate scope and stability issues are also addressed, alongside prospects for future research involving novel Schiff base frameworks and sustainable synthesis routes. This comprehensive overview serves as an essential resource for scientists exploring Schiff base complexes in bioinorganic and medicinal chemistry.

Keywords - Schiff bases; transition metal complexes; azomethine ligands; synthesis; biological activity; antimicrobial agents; catalysis; coordination chemistry; structure-activity relationship; green chemistry.

INTRODUCTION

Schiff bases, first described in 1864 by Hugo Schiff, are organic compounds characterized by the azomethine group (-C=N-), generally formed through a condensation reaction between primary amines and aldehydes or ketones under mild conditions. Owing to their synthetic accessibility and structural versatility, Schiff bases are widely employed as ligands in coordination chemistry, forming stable complexes with transition metals [2][3]. These complexes demonstrate variable coordination geometries and exhibit fascinating physicochemical properties, which include biological activities such as antimicrobial, antifungal, and antioxidant effects, along with catalytic capabilities in organic transformations [1][4].

The early 21st century brought remarkable advancements in Schiff base chemistry, particularly from 2000 to 2013, with numerous studies focusing on synthesizing transition metal complexes based on Schiff bases and exploring their potential in medicine and catalysis. The prevalent metals involved include cobalt, nickel, copper, palladium, manganese, and iron, whose coordination environments significantly influence the bioactivity and catalytic performance of resultant complexes [3][5].

This review synthesizes the key developments in this field during the past 14 years, highlighting synthetic strategies, structural elucidation, biological assays, and catalytic applications. Challenges and potential future directions are also outlined.

Synthetic Strategies for Schiff Base Ligands and Their Metal Complexes

The synthesis of Schiff bases commonly follows a straightforward condensation of an aromatic or aliphatic aldehyde with a primary amine, yielding ligands featuring the azomethine group essential for metal coordination. Aromatic aldehydes are often favored due to the enhanced stability of the resulting ligands derived from conjugation, whereas aliphatic Schiff bases frequently exhibit issues of polymerization and instability under ambient conditions [2][3].

Metal complexes are typically synthesized by reacting these ligands with metal salts such as chlorides, nitrates, or acetates of Ni(II), Co(II), Cu(II), Pd(II), Mn(II), and Fe(II) under reflux in polar solvents like ethanol, methanol, or dimethyl sulfoxide (DMSO). The stoichiometric ratios are usually 1:1 or 1:2 metal-to-ligand, depending on the denticity of the Schiff base ligand and the metal coordination preference [1][4].
Microwave-assisted synthesis has emerged as a powerful alternative to classical reflux, reducing reaction times and

© 2015 IJSRET

improving yields, while greener methodologies are being actively explored to address environmental concerns [5].

Structural Characterization

The complexes' structures are primarily elucidated through spectroscopic techniques complemented by elemental and thermal analyses. Key characterization methods include:

- Fourier Transform Infrared (FT-IR) Spectroscopy: Shifts in the characteristic azomethine (C=N) stretching frequency upon complexation provide evidence for ligand coordination through nitrogen atoms. Additional bands for metal-oxygen and metal-nitrogen bonds confirm bonding modes [3][4].
- Nuclear Magnetic Resonance (NMR) Spectroscopy: ^1H and ^13C NMR assist in confirming ligand structure and substitution patterns, though paramagnetic metal complexes often preclude detailed NMR studies of the complex [2].
- Ultraviolet-Visible (UV-Vis) Spectroscopy: Indicates d–d transitions and charge-transfer bands, providing insight into the coordination environment and geometry around the metal centers [4].
- Mass Spectrometry: Confirms molecular ion peaks corresponding to the expected molecular weights, affirming complex formation and fragmentation patterns [1].
- X-ray Crystallography: When crystallographic data is available, it definitively establishes coordination geometry, bond lengths, and packing modes critical for understanding structure-activity relationships [1].

Complexes of transition metals often adopt octahedral, square planar, or tetrahedral geometries, dictated by the metal ion and ligand denticity. For instance, Ni(II) and Co(II) tend to form octahedral complexes with tetradentate Schiff bases, while Cu(II) frequently prefers distorted square planar environments [3].

II. BIOLOGICAL ACTIVITIES

Transition metal Schiff base complexes exhibit enhanced biological activities compared to free Schiff bases due to the increased lipophilicity and altered electronic properties bestowed by metal coordination. The following summarizes key bioactivities reported between 2000 and 2013:

Antimicrobial and Antifungal Activity

Numerous studies demonstrate that Schiff base complexes effectively inhibit pathogenic bacteria such as Staphylococcus aureus, Escherichia coli, and Pseudomonas aeruginosa, along with fungi like Candida albicans [2][4]. Metal coordination enhances cell membrane penetration, disrupts microbial enzymes, and interferes with cellular DNA [3]. Complexes of Co(II), Ni(II), and Cu(II) have been widely investigated, with some showing superior activity over standard drugs in in vitro assays [4].

Antioxidant and Anticancer Activities

Certain Schiff base complexes show promising antioxidant properties, scavenging free radicals and protecting biomolecules from oxidative stress [3]. Complexes involving Mn(II) and Fe(II) have attracted attention for their redox activity linked to anticancer behavior, inducing apoptosis in malignant cells [1][5].

Other Biological Effects

Additional applications include antiviral, anti-inflammatory, and enzyme inhibitory activities, expanding the therapeutic potential of Schiff base metal complexes [2][5].

Catalytic Applications

Transition metal Schiff base complexes serve as efficient catalysts in various organic reactions, leveraging their redox flexibility and coordination geometries:

- Oxidation Reactions: Complexes of chromium, manganese, copper, and ruthenium effectively catalyze epoxidations and oxidation of olefins and alcohols [1].
- Polymerization: Organometallic cobalt complexes with tridentate Schiff bases have been utilized as initiators for emulsion polymerization of vinyl and dienyl monomers, accelerating polymer growth [2].
- Hydrogenation and Carbonylation: Selective homogeneous catalysts have been developed using Pd(II) and Ni(II) Schiff base complexes, facilitating these industrially relevant reactions with good yield and selectivity [1].

The catalytic performance is strongly influenced by metal identity, ligand electronic properties, and coordination environment, emphasizing the need for tailored ligand design [4].

Challenges and Future Perspectives

Despite promising advances, several challenges persist:

- Ligand Stability: Aliphatic Schiff bases suffer from polymerization and reduced shelf life, limiting their practical utility [2].
- Selective Activity: Enhancement of target selectivity and minimization of cytotoxicity require deeper mechanistic insights [3].
- Green Chemistry: Traditional synthetic procedures often involve toxic solvents and prolonged reflux, calling for

© 2015 IJSRET

greener approaches such as solvent-free, microwave, or ultrasound-assisted synthesis [5].

Emerging research seeks to integrate Schiff base complexes with nanotechnology platforms to enhance biological targeting and catalytic efficiency, alongside expanding water-compatible and biodegradable ligand frameworks.

III. CONCLUSION

The period from 2000 to 2013 has seen substantial progress in Schiff base coordination chemistry, resulting in diverse transition metal complexes with valuable biological and catalytic functions. The azomethine linkage and ligand design enable versatile and stable coordination, while metal complexation substantially boosts bioactivity and catalytic efficacy. Continued research addressing synthesis sustainability, stability improvement, and mechanistic understanding will further advance their practical application in medicinal chemistry and industrial catalysis.

REFERENCES

1. A. A. Galal et al., "Metal Complexes of Macrocyclic Schiff-Base Ligand: Preparation and Catalytic Applications," Journal of Coordination Chemistry, 2013.
2. Prakash, A., & Adhikari, D. (2011). Application of Schiff bases and their metal complexes-A Review. Int. J. Chem. Tech. Res, 3(4), 1891-1896.
3. Katwal, R., Kaur, H., & Kapur, B. K. (2013). Applications of copper—Schiff's base complexes: a review. Sci Rev Chem Commun, 3, 1-15.
4. S. M. Shehata et al., "Biological Activities of Schiff Bases and Their Complexes," Scientific Research Publishing Inc., 2013.
5. Prakash, A., & Adhikari, D. (2011). Application of Schiff bases and their metal complexes-A Review. Int. J. Chem. Tech. Res, 3(4), 1891-1896.

© 2015 IJSRET