International Journal of Scientific Research & Engineering Trends
Volume 2, Issue 4, Jul-Aug-2016, ISSN (Online): 2395-566X

Atmospheric Chemistry of Greenhouse Gases and Their Role in Global Warming

Dr. Sarika Sharma
Department of Chemistry
Government Degree College, Budaun, Uttar Pradesh, India.

Abstract — Atmospheric chemistry plays an important role in the global climate system as greenhouse gases (GHGs) are involved in the Earth's climate system, radiation, and atmosphere. GHGs such as carbon dioxide (CO_2), methane (CH_4), nitrous oxide (N_2O), and halogenated compounds absorb infrared light and emit it in the atmosphere of Earth as greenhouse gases, and this is associated with the greenhouse effect. The heat in the lower atmosphere is retained, and global warming and the surface temperature of the Earth are increasing. As such, the chemistry of greenhouse gases depends on the concentration of atmospheric gases as well as their chemical composition, reactivity, lifetime, and interaction with solar and terrestrial radiation (e.g., photochemical reactions, oxidation processes, gas-aerosol interaction). For example, methane oxidation and nitrogen oxide cycles play an important role in ozone production and secondary radiative forcing, so that the chemistry of atmospheric chemistry and climate are interrelated. Since the 20th century, anthropogenic activities such as combustion of fossil fuels, industrial pollution, deforestation, and agricultural processes have increased the GHG levels in our atmosphere, thus adding to the natural greenhouse effect. However, CO_2 is the most important greenhouse gas present now, but it is not the only one that is responsible for warming, and other gases such as CH_4 and N_2O are essential in the global warming process as well. Atmospheric chemistry reveals that the greenhouse effect is not only dependent on CO_2, but many interacting gases are involved in the climate processes. Recent studies have also shown that changes in the composition of the atmosphere can lead to severe weather events, radiative forcing, and climate feedback loops, and the consequences can be dramatic for global warming. In any system for climate change, the interplay of greenhouse gases, aerosols, and chemical reactions in the atmosphere should be taken into account. From a global perspective, understanding the chemistry and nature of greenhouse gases is necessary to understand what is driving us toward global warming. The chemical properties and interactions of these gases are also useful in understanding how climate change must be countered in the long run and how to identify solutions to this problem for climate policy.

Keywords: Atmospheric Chemistry, Greenhouse Gases (GHGs), Global Climate System, Greenhouse Effect, Global Warming, Carbon Dioxide (CO_2), Methane (CH_4), Nitrous Oxide (N_2O), Radiative Forcing, Photochemical Reactions, Atmospheric Oxidation Processes, Gas–Aerosol Interactions.

INTRODUCTION

The atmosphere plays a key role in the Earth's ecosystem by controlling the composition, transformation and radiative behaviour of gases residing in the atmosphere. Among these gases, greenhouse gases (GHGs) can absorb infrared and emit radiational energy which is directly related to the power balance in our planet. The greenhouse effect is a natural and important phenomenon. When a solar beam of shortwave light comes into a vacuum of Earth (in the atmosphere) then photons of this short wave radiation are attracted from the outer surface of Earth and then absorbed and emitted in the atmosphere, these photons are transferred back into the atmosphere on a long wave basis (and not yet absorbed). Greenhouse gases can absorb these photons so that they are in all directions and transmit them back to a warm climate. From a chemical perspective, greenhouse gases in their molecular structure (in terms of vibrational properties) control their response to infrared radiation. Unlike N_2 and O_2 which are very opaque to infrared radiation, we often hear about carbon dioxide (CO_2), methane (CH_4), nitrous oxide (N_2O) and chlorofluorocarbons (CFCs). These substances are very transparent and tend to have two different atomic structures to filter out temperature and, as the main component of radiative forcing (and hence, global warming). The chemistry of gases is further affected by photochemical and oxidation reactions as well as in the troposphere and stratosphere. Methane undergoes oxidation reactions to raise ozone and water vapor, both of which are greenhouse gases, for instance. Nitrogen oxides and volatile organic compounds (VOCs) also occur in photochemical cycles that also contribute to ozone and chemistry elsewhere as well as different atmospheric processes. Thus global warming is thus driven by the complex chemical interactions of the gases on the atmosphere but is a multi-component process.

The anthropogenic combustion of fossil fuels, industrialisation, deforestation and intensive agriculture in the past decades has altered the natural equilibrium between atmospheric gases, and

© 2016 IJSRET

the concentration of such gases in the atmosphere has increased significantly since the pre-industrial era, and this in turn has enhanced the greenhouse effect to a level that is able to lead to real global temperatures. In fact, the current research results suggest that the mean surface temperature has actually risen by about 1.1 °C above pre-industrial times and this is not expected to stop if greenhouse gases are not inhibited. In addition, atmospheric chemistry shows that the impact of greenhouse gases goes beyond carbon dioxide only. CO_2 is the most abundant anthropogenic greenhouse gas and so other gases (methane and nitrous oxide) are also significant warming agents on a per molecule basis. Halogenated gases, despite their present concentration, also play their part, as there is too much (especially radiative) long life of this gas in the atmosphere. What we are trying to see with global warming from a multi-gas perspective. In addition, the interaction of greenhouse gases with aerosols, clouds and atmospheric circulation introduces new layers of complexity of climate dynamics.

Depending on the composition and distribution of the air component (of the atmosphere) the warming will be enhanced or decreased because feedback mechanisms such as water vapor amplification and ice-albedo changes can be very efficient at accelerating climate change. This all shows us that atmospheric chemistry is present in physical and climatic systems and that accurate climate modeling or prediction is required. The atmospheric chemistry of greenhouse gases should be understood so that our understanding in the process can be useful in estimating its effects with regard to global warming. Not only does it provide the most fundamental knowledge of radiative forcing and climate feedbacks but also is the basis for mitigation approaches such as emission control, carbon capture and sustainable environmental management strategies.

II. MOLECULAR MECHANISMS OF HEAT TRAPPING

The ability of greenhouse gases (GHG) to trap heat in the atmosphere is very much determined by their molecular structure and interaction with the IR radiation. For a gas to be "greenhouse active," it either needs to have a permanent dipole moment or to have a temporary dipole in vibrational motion. If it does exist, a molecule would have its moment with the longwave light scattered by the Earth's surface. This is the only thing which is the greenhouse effect. The symmetry and vibration of the molecular system of homonuclear diatomic molecules such as nitrogen (N_2) and oxygen (O_2), which make up about 99% of Earth's atmosphere, are symmetrical and non-polar. There's no dipole movement and therefore no contact with IR radiation. And thus, although they are very high in quantity, they do not contribute to heat trapping or even global warming. However, carbon dioxide (CO_2), methane (CH_4), nitrous oxide (N_2O), and water vapor (H_2O) have asymmetric structures that change the dipole moment and are very good agents for IR absorption.

And of particular interest is the spectrum of absorption bands that are related to infrared radiation absorbed by greenhouse gases. As it turns out, all gases have their own vibrational frequency that shows which part of the electromagnetic spectrum they can absorb. For example, CO_2 absorbs the strongest IR radiation at ~15 μm, and water vapor has a much wider spectrum (at many wavelengths). Their absorption bands together determine the radiative forcing potential of each gas. Of particular interest is the atmospheric window (8-12 μm) where the emitted infrared radiation from the Earth can easily escape with minimal absorption.

Gases that absorb infrared radiation in that region are important because they are able to block one of the main paths for heat loss into space from the Earth. Methane (CH_4) and ozone (O_3) absorb radiation in the window region with very high warming potential even in the lower atmosphere. Apart from absorption, greenhouse gases also re-emit their absorbed energy into the lower atmosphere in numerous ways from the atmosphere even as they approach the surface of the Earth. The constant loop of absorption and re-radiation, thus, gives greenhouse gases an overall warming effect as the temperature remains in the lower atmosphere and thus a net warming effect.

The efficiency of this process depends on a number of variables such as molecular complexity, atmospheric concentration, and lifetime of the gas. In the end, molecular mechanisms of heat trapping show the importance of chemical structure and vibrational dynamics in global warming of different gases. This will be useful in the assessment of the impact of different greenhouse gases on global warming and in the design of solutions to their impact. Greenhouse gases vary in source, chemical properties, lifetime in atmosphere, and impact on global warming. We need to identify them and understand their individual impacts and plan for their mitigation to address climate change.

Carbon Dioxide (CO_2)

Although CO_2 is the most abundant anthropogenic greenhouse gas, mainly due to its high concentration and long atmospheric lifetime it is the main agent of global warming because it is the primary driver of the global warming process due to its high concentration and long residence. Most of the carbon dioxide is produced by burning fossil fuels (coal, oil, gas),

© 2016 IJSRET

deforestation, cement production, and natural processes such as volcanic activity and respiration, due to which it is easy to release carbon dioxide to the atmosphere. CO_2 is relatively inert and stable in the troposphere from a chemistry perspective because of its long residence time and does not undergo chemical change in an environment of normal atmospheric conditions because most of the time it is in a long residence period. The concentration of CO_2 is thus controlled by the global carbon cycle, in which photosynthesis removes CO_2 from the atmosphere (the photosynthesis and respiration of CO_2 and ocean absorption and respiration is the process for respiration in the troposphere. The oceans are the primary sink that pulls CO_2 out of the atmosphere and dissolves it into bicarbonate and carbonate ions. While CO_2 is less efficient in radiative production than other gases, its abundance and persistence over decades (from centuries to millennia) ultimately render it the primary source of climate change. Its duration and cumulative impact is one of the factors of long-term global warming.

Methane (CH_4)

Methane (CH_4) is a rather powerful greenhouse gas and has a much higher heat-trapping capacity than CO_2. Agricultural processes such as enteric fermentation in ruminants and rice paddies, landfills, wetlands, and leaks from natural gas extraction and distribution systems are the main sources of methane. Methane is chemically reactive and is most oxidized by the hydroxyl radical (·OH) (called the "atmospheric detergent"). The first reaction is:

$$CH\,4 + \cdot OH \rightarrow \cdot CH3 + H2O$$

This reaction is followed by oxidation (decrease) of methane, resulting in CO_2 and water vapor (H_2O). This also occurs in the stratosphere, which can lead to stratospheric water vapor formation as a greenhouse gas. Methane has a much shorter atmospheric lifetime (about 10 to 12 years) than CO_2, and the Global Warming Potential over 100 years (GWP-100) is about 28 times higher. Indeed, CH_4 is a key part of short-term climate mitigation efforts as the reduction of methane emissions can bring about immediate climate changes.

Nitrous Oxide (N_2O)

Nitrous oxide (N_2O) is another prominent greenhouse gas which is mainly produced by agricultural processes (especially nitrogen-based synthetic fertilizers) and by industrial processes and burning of biomass. In terms of chemistry, N_2O is very stable in the troposphere, and has low reactions at lower levels of atmosphere. Due to its high stability, it is known to last for over 100–120 years in the atmosphere. It finally reaches the stratosphere, where it either decomposes photochemically or reacts with excited oxygen atoms:

$$N2O + h\nu \rightarrow N2 + O\mathrm{N2}$$

O+hν→N2+O

Moreover, it plays a key role in stratospheric ozone depletion. It is currently the most ozone-depleting substance of the 21st century, and is linked to climate change and ozone chemistry. At a GWP-100 of about 273, nitrous oxide is far more efficient at trapping heat than CO_2 on a per molecule basis. Its role in both global warming and ozone depletion makes it of particular importance in atmospheric chemistry studies.

Tropospheric Ozone (O_3)

Tropospheric ozone (O_3) is a secondary greenhouse gas and is not emitted directly into the atmosphere but rather produced by complex photochemical reactions of nitrogen oxides (NO_x) and volatile organic compounds (VOCs) in sunlight. The ozone is formed along a series of reactions due to solar radiation that breaks down nitrogen dioxide (NO_2) and gives rise to ozone. Tropospheric ozone is a pollutant and greenhouse gas. Ozone is highly reactive and has a very short atmospheric lifetime (hours to weeks) but also has radiative forcing as it can absorb infrared light very well. It is also related to photochemical smog and can have negative effects on human health, vegetation, and ecosystems.

IV. COMPARATIVE ANALYSIS: GWP AND RADIATIVE FORCING

The climatic impact of greenhouse gases (GHGs) is measured based on two main factors: Radiative forcing (RF) and Global Warming Potential (GWP). Radiative forcing is the change in Earth's energy balance in watts per square meter (W/m^2) due to the change in the atmospheric composition since the pre-industrial era. Positive radiative forcing is a net gain of energy in the earth-atmosphere system, leading to warming and negative forcing is cooling. GWP is the relative effect of a greenhouse gas in terms of heat-trapping capacity of CO_2 over a given period of time (usually 100 years). From the atmospheric chemistry perspective, RF and GWP are influenced by many interrelated factors including the infrared absorption efficiency, the atmospheric lifetime, chemical reactivity and interaction with other atmospheric constituents. These factors all contribute to how long a gas remains in the atmosphere and how effectively it contributes to global warming.

Carbon dioxide (CO_2) is the reference gas with a GWP value of 1. Its radiative efficiency per molecule is relatively low, but its long atmospheric lifetime -- from centuries to millennia -- and high concentration mean that it is the dominant factor of radiative forcing. Its removal is limited by slow processes like

© 2016 IJSRET

oceanic dissolution and uptake by the terrestrial biosphere and its elimination from the atmosphere is slow (Table 1).

Methane (CH_4), by contrast, has a much shorter atmospheric lifetime of about 12 years and has a much higher GWP, typically in the range of 27 -30 over a 100 year period. Methane is also highly effective in absorbing infrared light in the troposphere due to its oxidation by hydroxyl radicals (·OH). Methane chemistry is also related to wider atmospheric oxidation. Despite its shorter lifetime, methane is very strong in radiative forcing near-term due to its high potency (Table 1). Nitrous oxide (N_2O) is another important greenhouse gas and has a long atmospheric lifetime of 121 years with a very high GWP of ~273. Its stability in the troposphere allows it to persist for many years before reaching the stratosphere where it is photolyzed or reacts with excited oxygen atoms. Besides its strong radiative contribution, N_2O is also an ozone-depleting substance and is in fact also crucial for atmospheric chemistry and climate change (Table 1).

Chlorofluorocarbons (CFCs), such as CFC-11, are among the most potent greenhouse gases, with extremely high GWP values (around 4,660) despite the relatively lower atmospheric concentrations. Their long life (~45 years) allows them to accumulate and have a strong warming effect. CFCs are mainly removed by photodissociation in the stratosphere, which also releases chlorine radicals responsible for ozone layer depletion. Their high radiative efficiency is because of their absorption in the atmospheric window region, which makes their warming potential much greater (Table 1).

Table 1: Comparative analysis of major greenhouse gases based on atmospheric lifetime, global warming potential (GWP-100), and dominant chemical removal pathways influencing their radiative forcing and climate impact.

Gas	Lifetime (Years)	GWP (100-year)	Major Chemical Sink
CO_2	Variable (100–1000+)	1	Ocean/Terrestrial Biosphere
CH_4	~12	27–30	Tropospheric $\cdot OH$
N_2O	~121	273	Stratospheric Photolysis
$CFC-11$	~45	4,660	Photodissociation

A comparison of these gases demonstrates that the impact of these gases is not a single parameter but rather a combination of concentration, lifetime, and radiative efficiency. CO_2 is the main cause of climate change for the long-term, but CH_4 and N_2O are also the major contributors to warming in short- and medium-term due to their higher GWP. Synthetic gases such as CFCs, which are present in trace quantities, have a significant impact as they are very radiative gases.

V. FEEDBACK LOOPS AND FUTURE TRENDS

A crucial aspect of atmospheric chemistry in the context of global warming is climate feedback loops which may help to amplify or, in some cases, moderate the effects of greenhouse gases (GHGs). In particular, the water vapor feedback effect is the most important positive feedback on the climate system of Earth. As carbon dioxide (CO_2) concentrations increase, the radiative forcing increases and the global temperature increases. The warmer air is more able to hold water (so much more water evaporates from oceans, soils and plants). Because water vapor is a very effective greenhouse gas that has a very high infrared absorption at all wavelengths, this will only increase the heat trapping. In other words, it will only be the initial warming from CO_2 that will be amplified by further warming from the increased water vapor. Atmospheric studies have shown that this feedback mechanism may double the warming from CO_2 and is therefore one of the key contributors to climate change sensitivity. Besides the water vapor feedback mechanisms, biogeochemical feedback mechanisms are increasingly being identified as a crucial driver of climate change in the future.

Recent observational and model studies (2024-2026) indicate that CH_4 emissions are being affected more by natural processes in nature such as wetlands. As global temperatures rise, microbial activity in waterlogged soils increases and organic matter is slowly anaerobically decomposing, accelerating the production of methane. This is a classic positive climate feedback in which warming increases the release of additional greenhouse gases that will lead to greater warming. Permafrost regions are another positive feedback system. As the temperature increases, frozen soils begin to thaw and organic carbon that is buried beneath them is released in the form of CO_2 and CH_4. This not only increases atmospheric greenhouse gas concentrations but makes regional hydrology and ecosystem dynamics change. Similarly, warming oceans can reduce their capacity to absorb CO_2 and thus weaken one of the planet's major carbon sinks and accelerate atmospheric accumulation.

© 2016 IJSRET

Atmospheric chemistry also sheds light on chemical feedbacks between reactive species such as hydroxyl radicals (·OH) and methane which influence the lifetime of methane. However, the amount of these molecules in the atmosphere will also be affected by changes in atmospheric composition, temperature and humidity and thus may affect the rate at which methane is removed from the atmosphere. If the amount of ·OH is reduced, the methane lifetime will be longer, hence its warming will be higher. Anthropogenic and non-anthropogenic sources of greenhouse gases and climate change will likely be a key component of global warming in the future. Climate models are now incorporating these feedback mechanisms in the context of climate change to enhance their predictions and emphasize that the Earth system responds in a nonlinear and interconnected manner. The increasing importance of natural emissions (driven by warming itself) suggests that the mitigation strategy should not only focus on human-induced emissions, but also on feedback-driven increases in greenhouse gases as well.

VI. CONCLUSION

The atmospheric chemistry of greenhouse gases clearly demonstrates that global warming is far more complex than a simple increase in CO_2 concentration. Rather it is the result of a complex interplay of molecular properties, chemical reactivity, atmospheric lifetimes, and feedback mechanisms. The ability of different gases to trap heat is also dependent upon their molecular structure and vibrational properties that determine their interaction with infrared radiation. At the same time, their persistence in the atmosphere (years to millennia) is a factor in how long they affect the Earth's radiative balance. These factors, combined with the chemical changes and interactions in the atmosphere, are connected to climate change. Water vapor amplification and methane release from natural sources in the world also have feedback loops in which to play out. Such feedbacks can significantly increase the initial warming due to greenhouse gases and lead to nonlinear and potentially faster climate changes.

The increase in natural emissions, stimulated by rising temperatures, shows that the climate system is not static but dynamic and self-reinforcing. In this context, climate mitigation strategies need to be a dual and integrated approach. On the one hand, reducing long-lived greenhouse gases such as CO_2 is central to controlling long-term global temperature rise as their cumulative impact lasts for centuries. On the other hand, targeting short-lived but extremely strong climate forcers such as methane (CH_4) can give the climate a rapid benefit by reducing the rate of warming in the short run. This dual approach is crucial for maintaining the climate system in the long term and for preventing tipping points in the short term. Additionally, a full understanding of atmospheric chemistry has great implications for multi-gas mitigation policies, better emission inventories, and better climate models with chemical interactions and feedback. It is also an example of an interdisciplinary approach to chemistry, environmental science, and policy-making.

Future Perspectives

The future of the atmospheric chemistry of greenhouse gases research is on the way to a deeper understanding of chemical processes, climate dynamics and human activities in the Earth. As global warming continues to grow, future studies will increasingly look at high-resolution monitoring, advanced modeling, and novel mitigation technologies to better understand and address climate change. The most important direction is to increase the monitoring of atmospheric instruments for global warming, with satellite-based remote sensing and ground-based monitoring. The latter will allow for a better understanding of greenhouse gas concentration and emission and temperature variations at regional and global scales. Better observations and data will also improve climate models and improve estimates of future warming scenarios.

One of the next important areas is atmospheric chemistry–climate models that incorporate complex feedback mechanisms like water vapor amplification, methane release from wetlands and permafrost, and changes in the ability of the air to oxidize. Chemical kinetics and climate dynamics can be combined to better simulate greenhouse gases in different emissions pathways and environmental conditions. In mitigation efforts in the future we will have to focus on multi-gas and multi-scale reduction. The long-term reduction of CO_2 emissions is crucial but shorter-lived climate pollutants like CH_4, tropospheric ozone (O_3) and hydrofluorocarbons (HFCs) will be in increasing demand. Rapid reductions of these gases may generate immediate climate change benefits and help slow the speed of warming in the short term.

Technological advances are also expected to help drive down emissions. New technologies such as carbon capture, utilization and storage (CCUS), direct air capture (DAC) and nature-based solutions (e.g., afforestation, soil carbon sequestration) are expected to be key to reduce emissions and we are also optimistic that green chemistry and sustainable industries will help reduce greenhouse gas release at the source and carbon dioxide production. And artificial intelligence and machine learning techniques are also well-suited to climate science. Such tools can examine large data sets and make predictions, and find patterns in atmospheric chemistry that are not easily captured by traditional methods. This will greatly enhance our

© 2016 IJSRET

ability to predict climate trends and develop effective interventions in the future. In future perspectives we must also consider the interplay between policy integration and global collaboration. Solutions to global warming need a comprehensive international approach in line with scientific knowledge of atmospheric chemistry. Policies that aim to reduce emissions, scale up technology and create sustainable development to sustain long-term climate stability need to be consistent with scientific understanding.

REFERENCES

1. Rypdal, K., & Winiwarter, W. (2001). Uncertainties in greenhouse gas emission inventories—evaluation, comparability and implications. Environmental Science & Policy, 4(2-3), 107-116.
2. Keller, S. (2010). Scientization: putting global climate change on the scientific agenda and the role of the IPCC. Poiesis & Praxis, 7(3), 197-209.
3. Jacob, D. J. (1999). Introduction to atmospheric chemistry. Princeton University Press.
4. Zellner, R. (2000). John H. Seinfeld and Spyros N. Pandis: Atmospheric chemistry and physics, from air pollution to climate change.
5. Finlayson-Pitts, B. J., & Pitts, J. N. (2000). Chemistry of the upper and lower atmosphere. Academic Press.
6. Wallace, J. M., & Hobbs, P. V. (2006). Atmospheric science: An introductory survey (2nd ed.). Academic Press.
7. Lelieveld, J., Dentener, F. J., Peters, W., & Krol, M. C. (2004). On the role of hydroxyl radicals in methane oxidation. Nature, 452, 737–740.
8. Ehhalt, D., & Prather, M. (2001). Atmospheric chemistry and greenhouse gases. In J. T. Houghton et al. (Eds.), Climate change 2001: The scientific basis. Cambridge University Press.
9. Forster, P., et al. (2007). Changes in atmospheric constituents and radiative forcing. In Climate change 2007: The physical science basis. Cambridge University Press.
10. Myhre, G., et al. (2013). Anthropogenic and natural radiative forcing. In Climate change 2013: The physical science basis. Cambridge University Press.
11. Ramaswamy, V., et al. (2001). Radiative forcing of climate change. Reviews of Geophysics, 39(1), 71–122.
12. Prather, M. J., et al. (2012). Measuring and modeling the lifetime of nitrous oxide. Atmospheric Chemistry and Physics, 12, 5963–5990.
13. Monks, P. S., et al. (2009). Atmospheric composition change—global and regional air quality. Atmospheric Environment, 43(33), 5268–5350.
14. Crutzen, P. J. (1995). Ozone in the troposphere and stratosphere. Annual Review of Energy and the Environment, 20, 137–164.
15. Ravishankara, A. R., Daniel, J. S., & Portmann, R. W. (2009). Nitrous oxide as the dominant ozone-depleting substance. Science, 326(5949), 123–125.
16. Cicerone, R. J., & Oremland, R. S. (1988). Biogeochemical aspects of atmospheric methane. Global Biogeochemical Cycles, 2(4), 299–327.
17. Houghton, J. T. (2002). The physics of atmospheres (3rd ed.). Cambridge University Press.
18. Kiehl, J. T., & Trenberth, K. E. (1997). Earth's annual global mean energy budget. Bulletin of the American Meteorological Society, 78(2), 197–208.
19. Lacis, A. A., Schmidt, G. A., Rind, D., & Ruedy, R. A. (2010). Atmospheric CO_2: Principal control knob governing Earth's temperature. Science, 330(6002), 356–359.
20. Byrne, B., & Goldblatt, C. (2014). Radiative forcing at high concentrations of well-mixed greenhouse gases. *Geophysical Research Letters*, *41*(1), 152-160.
21. Meinshausen, M., Meinshausen, N., Hare, W., Raper, S. C., Frieler, K., Knutti, R., & Allen, M. R. (2009). Greenhouse-gas emission targets for limiting global warming to 2 C. *Nature*, *458*(7242), 1158-1162.

© 2016 IJSRET